ELSEVIER

Carbohydrate Research 263 (1994) C15–C17

CARBOHYDRATE RESEARCH

# CONTENTS

The composition of reducing sugars in dimethyl sulfoxide solution
S.J. Angyal (Sydney, Australia) . . . . . 1

A quantitative approach to describe the overall motion of monosaccharides in solution. A $^{13}C$ NMR relaxation study of 1,6-anhydro-β-D-glucopyranose in dimethyl sulfoxide
P. Dais (Iraklion, Crete, Greece) . . . . . 13

Complexation of trivalent lanthanide cations by sugars and alditols in water: chromatography–calorimetry comparison
Y. Israëli, J.-P. Morel and N. Morel-Desrosiers (Aubière, France) . . . . . 25

Crystal and molecular structure of allyl *O*-(sodium 3-deoxy-α-D-*manno*-2-octulopyranosylonate)-(2 → 8)-*O*-(sodium 3-deoxy-α-D-*manno*-2-octulopyranosidonate)-monohydrate
V. Mikol (Basel, Switzerland), P. Kosma (Wien, Austria) and H. Brade (Borstel, Germany) . . 35

Synthesis of 4,6:2′,3′:4′,6′-tri-*O*-cyclohexylidene-α,α′-trehalose 2-palmitate: an intermediate for the synthesis of mycobacterial 2,3-di-*O*-acyl-α,α′-trehalose antigens
P.A. Wallace and D.E. Minnikin (Newcastle upon Tyne, United Kingdom) . . . . . 43

One-pot synthesis of 1,2-*O*-isopropylidene-α-D-xylofuranose
J. Moravcová, J. Čapková and J. Staněk (Prague, Czech Republic) . . . . . 61

Synthesis of α-D-galactopyranosyl-linked oligosaccharides containing the α-Gal → β-Gal → GlcNAc sequence employing methyl-2,3,4,6-tetra-*O*-(4-methoxybenzyl)-1-thio-β-D-galactopyranoside as an efficient glycosyl donor
G.V. Reddy, R.K. Jain, B.S. Bhatti and K.L. Matta (Buffalo, NY, USA) . . . . . 67

Structure elucidation of an acetylated saponin of *Blighia welwitschii* by NMR spectroscopy
A. Penders and C. Delaude (Liege, Belgium) . . . . . 79

Characterisation of an anti-ulcer pectic polysaccharide from leaves of *Panax ginseng* C.A. Meyer
H. Kiyohara, M. Hirano, X.-G. Wen, T. Matsumoto, X.-B. Sun and H. Yamada (Tokyo, Japan) . . . . . 89

Composition of the coagulant polysaccharide fraction from *Strychnos potatorum* seeds
M. Adinolfi, M.M. Corsaro, R. Lanzetta, M. Parrilli (Napoli, Italy), G. Folkard, W. Grant and J. Sutherland (Leicester, United Kingdom) . . . . . 103

An antitumor, branched (1 → 3)-β-D-glucan from a water extract of fruiting bodies of *Cryptoporus volvatus*
S. Kitamura, T. Hori, K. Kurita, K. Takeo (Kyoto, Japan), C. Hara (Gifu, Japan), W. Itoh, K. Tabata (Kobe, Japan), A. Elgasaeter and B.T. Stokke (Trondheim, Norway) . . . . . 111

Studies of the transglycosylation reaction catalysed by the decycling maltodextrinase of *Flavobacterium sp.* no. 92 with malto-oligosaccharides and cyclodextrins
H. Bender (Freiburg i.Br., FRG) . . . . . 123

Studies of the action pattern on potato starch of the decycling maltodextrinase from *Flavobacterium sp.* no. 92
H. Bender (Freiburg i.Br., FRG) . . . . . 137

*Notes*
The unusually stable crystal structure of *neo*-inositol
S.J. Angyal and D.C. Craig (Sydney, Australia) . . . . . 149

Application of the $^{29}$Si NMR 2D INEPT spin-flip *J*-resolved technique to the structural analysis of trimethylsilylated glycosides
M. Harket, B. De Jeso, J.-C. Lartigue, M. Petraud and M. Ratier (Talence, France) . . . . . 155

Solubility of lactulose trihydrate
T. Mizota, I. Suzawa, N. Seki, Y. Tamura and S. Shimamura (Kanagawa, Japan) . . . . . 163

Inhibition of bacterial and viral sialidases by 3-fluoro-*N*-acetylneuraminic acid
T. Hagiwara, I. Kijima-Suda (Saitama, Japan), T. Ido, H. Ohrui (Sendai, Japan) and K. Tomita (Saitama, Japan) . . . . . 167

*Regular papers*
Synthesis, NMR, and conformational studies of methyl α-D-mannopyranoside 2-, 3-, 4-, and 6-monophosphates
C. Bernlind, S. Oscarson and G. Widmalm (Stockholm, Sweden) . . . . . 173

Total synthesis of a modified ganglioside, de-*N*-acetyl $GM_2$
S. Fujita, M. Numata, M. Sugimoto, K. Tomita (Saitama, Japan) and T. Ogawa (Tokyo, Japan) . . . . . 181

The thermodynamic characteristics of the conformational transitions of native xanthan
L. Bezemer, M.E. Kuil and J.C. Leyte (Leiden, Netherlands) . . . . . 197

Structural studies of the *Escherichia coli* O90 O-antigen polysaccharide
S. Ratnayake, G. Widmalm (Stockholm, Sweden), A. Weintraub and E.C. Medina (Huddinge, Sweden) . . . . . 209

Structural comparison of the O6 specific polysaccharides from *E. coli* O6:K2:H1, *E. coli* O6:K13:H1, and *E. coli* O6:K54:H10
B. Jann (Freiburg, Germany), A.A. Shashkov (Moscow, Russian Federation), H. Kochanowski and K. Jann (Freiburg, Germany) . . . . . 217

Isolation and purification of feruloylated oligosaccharides from cell walls of sugar-beet pulp
M.-C. Ralet (Nantes, France), C.B. Faulds, G. Williamson (Norwich, United Kingdom) and J.-F. Thibault (Nantes, France) . . . . . 227

Structure identification of feruloylated oligosaccharides from sugar-beet pulp by NMR spectroscopy
I.J. Colquhoun (Norwich, United Kingdom), M.-C. Ralet, J.-F. Thibault (Nantes, France), C.B. Faulds and G. Williamson (Norwich, United Kingdom) . . . . . 243

Degradation of feruloylated oligosaccharides from sugar-beet pulp and wheat bran by ferulic acid esterases from *Aspergillus niger*
M.-C. Ralet (Nantes, France), C.B. Faulds, G. Williamson (Norwich, United Kingdom) and J.-F. Thibault (Nantes, France) . . . . . 257

Heparin-like compounds prepared by chemical modification of capsular polysaccharide from *E. coli* K5
B. Casu, G. Grazioli (Milan, Italy), N. Razi (Uppsala, Sweden), M. Guerrini, A. Naggi, G. Torri, P. Oreste, F. Tursi, G. Zoppetti (Milan, Italy) and U. Lindahl (Uppsala, Sweden) . . . . 271

Sequencing of xyloglucan oligosaccharides by partial Driselase digestion: the preparation and quantitative and qualitative analysis of two new tetrasaccharides
E.P. Lorences and S.C. Fry (Edinburgh, United Kingdom) . . . . . 285

*Notes*
Convenient synthetic approach towards C-3 modified methyl β-lactosides
Y. Nishida and J. Thiem (Hamburg, Germany) . . . . . 295

Sugar β-ketoesters: new chirons in the synthesis of 6-deoxyheptulosurono-7,4-lactones
D.D. Dhavale, N.N. Bhujbal, P. Joshi and S.G. Desai (Pune, India) . . . . . 303

Unsaturated ketonucleotides: synthesis of α and β anomers of 1-(2,3-dideoxy-6-*O*-diethoxyphosphoryl-D-*glycero*-hex-2-enopyranosyl-4-ulose)thymine
F. Leclercq and K. Antonakis (Villejuif, France) . . . . . 309

FTIR studies of sodium hyaluronate and its oligomers in the amorphous solid phase and in aqueous solution
R. Gilli (Trieste, Italy), M. Kacuráková, M. Mathlouthi (Reims, France), L. Navarini and S. Paoletti (Trieste, Italy) . . . . . 315

The structure of the *Citrobacter freundii* O8a,8b O-specific polysaccharide containing D-xylofuranose
N.A. Kocharova, Y.A. Knirel, A.S. Shashkov, N.K. Kochetkov, E.V. Kholodkova and E.S. Stanislavsky (Moscow, Russian Federation) . . . . . 327

*Preliminary communication*
Total synthesis of sulfated $Le^x$ pentaosyl ceramide
S. Numomura, M. Iida, M. Numata, M. Sugimoto (Saitama, Japan) and T. Ogawa (Tokyo, Japan) . . . . . C1

*Book review* . . . . . C7

*Author index* . . . . . C9

*Subject index* . . . . . C11

*Contents* . . . . . C15